

## *Pannaria hookeri* (lichenised Ascomycetes) – a remarkable new record for Australia

Gintaras Kantvilas<sup>1</sup> and Cécile Gueidan<sup>2</sup>

<sup>1</sup> Tasmanian Herbarium, Tasmanian Museum and Art Gallery, P.O. Box 5058, UTAS LPO, Sandy Bay, Tasmania 7005. Email: Gintaras.Kantvilas@tmag.tas.gov.au

<sup>2</sup> Australian National Herbarium, National Research Collections Australia, National Facilities and Collections, CSIRO, PO Box 1700, Canberra ACT 2601. Email: Cecile.Gueidan@csiro.au

### Abstract

*Pannaria hookeri* (Borrer ex Sm.) Nyl., a bipolar lichen, is recorded for the first time for Australia (south-west Tasmania) where it grew on alpine limestone outcrops. Its identity was confirmed by morphological, anatomical and DNA-sequence data.

**Keywords:** biodiversity, lichens, species discovery, Tasmania

### Introduction

Additions to the list of lichens recorded for Australia are made almost continuously, as demonstrated very well by McCarthy (2017), whose checklist for Australia requires regular updates. It is no exaggeration to observe that, in Australia, a survey of almost any area, or a revision of any taxonomic group, will reveal previously overlooked novelties. The Bush Blitz Programme, conducted by the Australian Biological Resources Study since 2010, has been especially successful in this regard. This paper reports a particularly interesting discovery, *Pannaria hookeri* (Borrer ex Sm.) Nyl., a first record for Australia. This bipolar species is widespread in arctic-alpine areas of the Northern Hemisphere (Jørgensen 1978; Thomson 1984) with scattered occurrences in the Antarctic (Øvstedal & Lewis Smith 2001), Mt Kenya in Africa (Jørgensen 2007), southern South America (Jørgensen 1978) and New Zealand (Galloway 1985). It was discovered in south-western Tasmania during a Bush Blitz Expedition in 2016.

## Methods

### Morphology, anatomy and chemistry

Morphological and anatomical investigations were undertaken on material collected in Tasmania and compared with reference herbarium material of *P. hookeri* as listed. Hand-cut sections were mounted in water, 10% KOH, 50% HNO<sub>3</sub> and Lactophenol Cotton Blue. Ascospore measurements are based on 75 observations and are presented in the format: least value–average–highest value; outlying values are given in brackets. Routine chemical analyses by thin-layer chromatography follow standard methods (Orange et al. 2001).

#### *Selected comparative specimens examined:*

**SWEDEN.** Torne Lappmark, Jukkasjärvi par., along river Lektajohka, c. 14 km ESE of Riksgränsen, 4.vii.2015. M. Westberg VAR 192 (S); Lycksele Lappmark: par. Tärna, Rivovardo, 800–1000 m alt., 1924, A.H. Magnusson 7922 (S). **NORWAY.** Opland: Nord-Fron hd, Sikilsdalslshö, c. 1620 m. 1620 m alt., 6.ix.1949, S. Ahlner (S). **AUSTRIA.** Nordtirol, Kluppescharte, 1960, H. Doppelbauer & J. Poelt (Lichenes Alpium 141) (MEL 1017675, S).

### DNA extraction, amplification and sequencing

Part of the specimen GK102/16 was sampled for DNA extraction using a clean pair of tweezers. The sampled material was transferred to an Eppendorf tube and the DNA extracted using a DNeasy® Plant Mini Kit (Qiagen, Hilden, Germany). The manufacturer's protocol was modified as follows: the material was ground in AP1 buffer using a sterile plastic pestle and then incubated with RNase A for an hour at 70°C. The lengths of all centrifuge steps were doubled and the DNA was recovered in 50 µl of AE buffer.

Two gene regions were amplified: the region including the 5.8S subunit of the nuclear ribosomal RNA gene and the internal transcribed spacers 1 and 2 (ITS) and the small subunit of the mitochondrial ribosomal RNA gene (mtSSU). The primers used were ITS1F (Gardes & Bruns 1993) and ITS4 (White et al. 1990) for ITS and mtSSU1 and mtSSU3R (Zoller et al. 1999) for mtSSU. DNA extracts were checked with a gel electrophoresis and for each sample the band intensity was used to choose the appropriate genomic DNA dilution for amplification.

One micro-litre (µl) of a 1, 1/10, or 1/100 dilution of genomic DNA was added to the following PCR mix: 2.5 µl PCR buffer 10 × NH<sub>4</sub> (Bioline, London, U. K.), 1.5 µl of MgCl<sub>2</sub> (50 mM), 0.5 µl dNTP (100 mM), 2.5 µl of BSA (10 mg/ml), 1 µl of each primer (10 µM), 0.25 µl DNA polymerase Bioline BioTaq (5 U µl<sup>-1</sup>), and water to a total volume of 25 µl. The PCR reactions were performed on a C1000Touch thermal cycler (Bio-Rad, Hercules, CA, USA). The PCR program for ITS was as follow: 5 min at 94°C, followed by 35 cycles of the three steps 1 min at 94°C (denaturation), 1 min at 53°C (annealing), 2 min at 72°C (extension), and a final extension time of 10 min at 72°C. For mtSSU, the program was: 3 min at 94°C, followed by 35 cycles of the three steps 1 min at 94°C, 1 min at 52°C, 1.5 min at 72°C, and a final extension time of 7 min at 72°C. Cloning was conducted on PCR products using a TOPO-TA Cloning kit (Invitrogen, Carlsbad, California), as instructed by the manufacturer. PCR product clean-up and sequencing were carried out by Macrogen (Seoul, South Korea) using BigDye chemistry and an ABI 3730xl sequencer (Applied Biosystems, Carlsbad, California).

### Phylogenetic analysis

New *Pannaria* sequences were edited using Sequencher v. 5.4.1 (Gene Codes Corporation, Ann Arbor, Michigan, USA). Published ITS and mtSSU *Pannaria* sequences were obtained from GenBank (<https://www.ncbi.nlm.nih.gov/>) and manual alignments of all sequences were done using Mesquite v. 3.04 (Maddison & Maddison 2011). The final alignments included 24 ingroup taxa from the genus *Pannaria* and three outgroup species (*Psoroma implexum*, *Staurolemma oculatum* and *S. omphalarioides*) were chosen, based on a previous phylogenetic study (Ekman et al. 2014). Taxon and gene sampling are shown in Table 1. Ambiguous regions were delimited as described in Lutzoni et al. (2000) and were excluded from the phylogenetic analyses. To test for congruence, each locus (ITS and mtSSU) was first subjected to a bootstrap analysis separately using maximum likelihood (ML) (RAxML VI-HPC v.8.2.9; Stamatakis et al. 2005, 2008), as implemented on the CIPRES Web Portal (<http://www.phylo.org>; Miller et al. 2010). A GTRCAT model was applied to the two markers. Support values were obtained using a fast bootstrap analysis of 1,000 pseudoreplicates. Resulting topologies

were compared for a potential conflict among loci using a 70% reciprocal bootstrap criterion (Mason-Gamer & Kellogg 1996). Because no conflict was detected, the two gene regions were concatenated and the combined dataset analysed using RAXML as described above. Trees were visualized in PAUP\* (Swofford 2002) and edited with Illustrator (Adobe Systems, San Jose, CA, USA). The dataset was deposited in TreeBase (22176).

Results

Phylogenetic results

The dataset included 27 taxa (24 ingroup and 3 outgroup taxa) and 1,114 characters (385 from ITS and 729 from mtSSU). The concatenated alignment had 228 distinct alignment patterns and a proportion of gaps and completely undetermined characters of 21.70%. The most likely tree is presented in Figure 1 with bootstrap support values. As in Ekman et al. (2014),

**Table 1.** Taxon and gene sampling for the phylogenetic analysis. Missing sequences are indicated by a dash. The accession numbers of the two newly generated sequences are highlighted in bold.

| Species   | Voucher                | ITS             | mtSSU           |
|---|------------------------|-----------------|-----------------|
| 1 <i>Pannaria andina</i>                          | Elvebakk 06-245        | GQ927268        | -               |
| 2 <i>Pannaria athraaphylla</i>                    | Passa 181              | EU885295        | EU885317        |
| 3 <i>Pannaria athraaphylla</i>                    | Passa 251              | EU885303        | EU885325        |
| 4 <i>Pannaria calaphylla</i>                      | Passa 101              | EU885296        | EU885318        |
| 5 <i>Pannaria canaplea</i>                        | Ekman 3188             | AF429281        | -               |
| 6 <i>Pannaria cantarta</i>                        | Passa 142              | EU885297        | EU885319        |
| 7 <i>Pannaria farinasa</i>                        | Passa 119              | EU885299        | EU885321        |
| 8 <i>Pannaria haakeri</i>                         | Jørgensen s.n.         | AF429282        | KC608083        |
| 9 <i>Pannaria haakeri</i>                         | GK102/16               | <b>MG786563</b> | <b>MG792317</b> |
| 10 <i>Pannaria immixta</i>                        | Elvebakk 02-352b       | -               | KC608084        |
| 11 <i>Pannaria insularis</i>                      | Kashiwadani 43760      | KC618716        | KC608085        |
| 12 <i>Pannaria leucasticta</i>                    | Hur 041227             | EU266107        | -               |
| 13 <i>Pannaria lurida</i> subsp. <i>lurida</i>    | Kashiwadani 43861      | -               | KC608086        |
| 14 <i>Pannaria lurida</i> subsp. <i>russellii</i> | Tønsberg 22565         | -               | KC608087        |
| 15 <i>Pannaria micraphyllizans</i>                | Passa 264              | EU885300        | EU885322        |
| 16 <i>Pannaria multifida</i>                      | Schumm & Frahm s.n.    | KC618717        | KC608088        |
| 17 <i>Pannaria pallida</i>                        | Passa 249              | EU885301        | EU885323        |
| 18 <i>Pannaria rubiginella</i>                    | Tønsberg 32508         | KC618718        | KC608089        |
| 19 <i>Pannaria rubiginella</i>                    | Thar 10050             | -               | GQ259037        |
| 20 <i>Pannaria rubiginasa</i>                     | Ananby 870/Purvis s.n. | AF429280        | AY340513        |
| 21 <i>Pannaria sphinctrina</i>                    | Passa 221              | EU885302        | EU885324        |
| 22 <i>Pannaria subfusca</i>                       | Tønsberg 33592         | KC618719        | -               |
| 23 <i>Pannaria tavaresii</i>                      | Schumm s.n.            | KC618720        | -               |
| 24 <i>Pannaria tavaresii</i>                      | Passa 122              | EU885294        | EU885316        |
| 25 <i>Psarama implexum</i>                        | Passa 84               | -               | EU885333        |
| 26 <i>Stauralemma aculatum</i>                    | Aptraat 55941          | KC618738        | GQ259045        |
| 27 <i>Stauralemma amphalariaides</i>              | Tibell s.n.            | KJ533487        | KJ533439        |




the genus *Pannaria* is supported as monophyletic (82% bootstrap support). The specimen of *Pannaria* collected in Tasmania (GK102/16) is highly supported as sister taxon to *Pannaria hookeri* (99% bootstrap support). The relatively short branch lengths between the Tasmanian and Northern Hemisphere specimens agree with their conspecific nature.

## Taxonomy

***Pannaria hookeri* (Borrer ex Sm.) Nyl., Mém. Soc. Natn. Sci. Nat. Cherbourg 5: 109 (1858)**

*Lichen hookeri* Borrer ex Sm., in J.E. Smith & J. Sowerby, *Engl. Bot.* 32: 2283 (1811).

**Type: SCOTLAND.** Ben Lawers and Meall Greigh, W. Borrer (lectotype, *fide* Jørgensen 1978: BM!).



**Figure 1.** Most likely tree showing the phylogenetic placement of the Tasmanian *Pannaria* specimen (GK102/16). The phylogeny was obtained using a combined ITS-mtSSU dataset and analysed with maximum likelihood using RAxML. Bootstrap support values are reported above the branches. The scale bar represents the number of nucleotide substitutions/site.

*Thallus* subcrustose to squamulose; squamules plane to convex, sometimes rather gnarled, 0.3–1 mm wide, 250–500 µm thick, dispersed or clustered and overlapping, tightly adnate or loosely attached over a black, effuse prothallus, irregularly rhomboidal and delimited by deep cracks in the centre of the thallus, ± effigurate and spatulate at the margins; upper surface pale brownish grey to smoky bluish grey, ± scabrid-maculate and faintly whitish striate, especially at the thallus margins; photobiont *Nostoc*, comprising single, roundish cells, 5–10 µm wide. *Apothecia* lecanorine, 0.4–1.3 mm diam., scattered, basally constricted when mature; disc dark brown to black, mostly plane but becoming convex and puckered with age; thalline margin mostly entire, crenulate, somewhat inrolled, in section 80–150 µm thick; proper margin prosoplectenchymatous, highly reduced to a dark brown band c. 10–20 µm wide between the hymenium and the thalline margin. *Hypothecium* colourless to pale yellowish, lacking photobiont cells. *Hymenium* (65–)70–90(–120) µm thick, colourless, I+ brown, K/I+ blue, overlain by an epithelial band of blue-green pigment, K+ intensifying greenish, N+ crimson. *Paraphyses* simple, 1.5–2.5 µm thick, with the terminal cell frequently enlarged to 4–5 µm and internally blue-green pigmented. *Asci* 8-spored but usually with a few ascospores deformed or aborted, (50–)55–65 x 14–20 µm, clavate, with a well developed, non-amyloid or very weakly amyloid tholus lacking any internal discernible structures, and an intensely amyloid, thin outer sheath. *Ascospores* broadly ellipsoid to ovate, smooth-walled, (10–)12–14.0–16(–18) x 6–7.6–9(–10) µm, when young with a distinct wall c. 1 µm thick and a large vacuole. *Pycnidia* not found. *Chemistry*: traces of pannarin sometimes found by t.l.c. but not detectable by spot tests. Figs 2–3.


**Specimen examined:** TASMANIA. North East Ridge of Mt Anne, at the western rim of Annakanada sinkhole, 42°55'57" 146°26'29"E, 1050 m alt., on sheltered limestone outcrops, 5.ii.2016, G. Kantvilas 102/16 (HO).

## Discussion

The above description is based solely on Tasmanian material but compares favourably with published descriptions (e.g. Jørgensen 1978; Thomson 1984; Galloway 1985; Stenroos et al. 2016), and with reference

herbarium material. Further confirmation of our determination was provided by molecular data.

Together with three other species that have highly restricted, circum-Antarctic distributions, *Pannaria hookeri* was placed in the subgenus *Chryopannaria* (Jørgensen 2000). However, subsequent phylogenetic research (Ekman & Jørgensen 2002; Ekman et al. 2014) did not support this classification and simply places *P. hookeri* within *Pannaria* s. str.



**Figure 2.** Morphology of *Pannaria hookeri* (GK 102/16). A. Habit; B. Detail of effigurate squamules with maculate, striate margins; C. Detail of apothecia. Scale = 2 mm




Lichens may be notoriously widespread, displaying cosmopolitan, bipolar, pan-temperate and pan-tropical distribution patterns, as well as having more localised ranges. However, applying names across wide geographic areas carries some risks, and there are numerous instances where a species perceived to be the same in both hemispheres has been subsequently found to comprise two distinct entities. For example, Australasian populations of what was once referred to as *Menegazzia terebrata* (Hoffm.) A.Massal. are now called *M. subpertusa* P.James & D.J.Galloway, and the lichen once called *Parmelia omphalodes* (L.) Ach. in Australasia is now correctly called *Notoparmelia signifera* (Nyl.) A.Crespo, Ferencova & Divakar. In this context, Robert Brown's list of lichens native to both Australia and Europe (Brown 1814) makes interesting reading.

In the case of *Pannaria hookeri*, the molecular investigation was critical. When first observed in the field and collected (by GK), the species was immediately recognised as truly novel for Tasmania. However, initial misinterpretation of some ambiguous apothecial characters led to assumptions that it was new to science and of uncertain generic affinity. It was the molecular

investigation, essentially to explore its generic relationships within the Pannariaceae, which redirected us towards *P. hookeri* and to making the necessary anatomical and morphological comparisons to confirm our identification.

### Ecology and distribution

The ecology of *Pannaria hookeri* is indeed unusual, for it occurs on alpine limestone, an extremely uncommon habitat in Tasmania. Interestingly, several European authors (e.g. Jørgensen 2000; Stenroos et al. 2016; James & Purvis 2009) likewise note its predilection for calcareous substrata. The species was part of a highly depauperate community on relatively sheltered aspects where lichens were patchy and extensive areas of bare rock prevailed. Other lichens recorded included *Baeomyces heteromorphus* Nyl. ex C.Bab. & Mitt., *Catillaria lenticularis* (Ach.) Th.Fr., *Lepraria vouauxii* (Hue) R.C.Harris, *Paraporphidia leptocarpa* (C.Bab. & Mitt.) Rambold & Hertel, *Placopsis brevilobata* (Zahlbr.) I.M.Lamb, *P. subcribellans* (I.M.Lamb) D.J.Galloway, *Porpidia crustulata* (Ach.) Hertel & Knoph, *P. umbonifera* (Müll.Arg.) Rambold, *Rhizocarpon petraeum* (Wulfen) A.Massal., *R. reductum* Th.Fr., *Staurothele succedens* (Rehm ex Arnold) Arnold, *Stereocaulon ramulosum* (Sw.) Räscher and a putative new species of *Trapelia* M.Choisy. Several of these taxa are themselves new or interesting records for Tasmania.



**Figure 3.** Anatomy of *Pannaria hookeri* (GK 102/16). Asci (amyloid parts stippled), paraphyses and ascospores. Note the typical *Pannaria*-type ascus with a well developed, non-amyloid tholus that lacks internal differentiation. Scale = 10  $\mu$ m

### Acknowledgements

A major part of this work was undertaken at the Swedish Museum of Natural History where G. Kantvilas was a guest researcher; Mats Wedin is thanked for his hospitality during that time. We thank Jean Jarman for the photographs and preparing all illustrations for publication. Tasmanian material studied was collected during a field survey co-funded by the Australian Biological Resources Study (ABRS) and BHP Billiton under the Bush Blitz Programme. Laboratory work was supported by an ABRS Tactical Taxonomy Grant to GK.

## References

- Brown, R. (1814). General remarks, geographical and systematical, on the botany of Terra Australis, in M. Flinders, *A Voyage to Terra Australis*, Volume 2, 533–613. G & W. Nicol, London.
- Ekman, S. and Jorgensen, P.M. (2002). Towards a molecular phylogeny for the lichen family Pannariaceae (Lecanorales, Ascomycota). *Canadian Journal of Botany* **80**, 625–634.
- Ekman, S., Wedin, M., Lindblom, L. and Jorgensen, P.M. (2014). Extended phylogeny and revised generic classification of the Pannariaceae (Peltigerales, Ascomycota). *Lichenologist* **46**, 627–656.
- Galloway, D.J. (1985). *Flora of New Zealand Lichens*. P.D. Hasselberg, Government Printer, Wellington.
- Gardes, M. and Bruns, T.D. (1993). ITS primers with enhanced specificity for Basidiomycetes: application to the identification of mycorrhizae and rusts. *Molecular Ecology* **2**, 113–118.
- James, P.W. and Purvis, O.W. (2009). Pannaria Delise ex Bory (1828), in C.W. Smith, A. Aptroot, B.J. Coppins, A. Fletcher, O.L. Gilbert, P.W. James and P.A. Wolseley (eds), *The Lichens of Great Britain and Ireland*, 650–651. British Lichen Society, London.
- Jørgensen, P.M. (1978). The lichen family Pannariaceae in Europe. *Opera Botanica* **45**, 1–123.
- Jørgensen, P.M. (2000). Studies in the lichen family Pannariaceae IX. A revision of Pannoria subg. Chryaponnorio. *Nova Hedwigia* **71**, 405–414.
- Jørgensen, P.M. (2007). Pannariaceae. *Nordic Lichen Flora* **3**, 96–112.
- Lutzoni, F., Wagner, P., Reeb, V. and Zoller, S. (2000). Integrating ambiguously aligned regions of DNA sequences in phylogenetic analyses without violating positional homology. *Systematic Biology* **49**, 628–651.
- Maddison, W.P. and Maddison, D.R. (2011). *Mesquite: a modular system for evolutionary analysis*, version 2.75.
- Mason-Gamer, R.J. & Kellogg, E.A. (1996). Testing for phylogenetic conflict among molecular data sets in the tribe Triticeae (Gramineae). *Systematic Biology* **45**, 524–545.
- McCarthy, P.M. (2017). *Checklist of the Lichens of Australia and its Island Territories*. Australian Biological Resources Study, Canberra. Version 12 April 2017. <http://www.anbg.gov.au/abrs/lichenlist/introduction.html>.
- Miller, M.A., Pfeiffer, W. and Schwartz, T. (2010). Creating the CIPRES Science Gateway for inference of large phylogenetic trees. 2010 Gateway Computing Environments Workshop (GCE), New Orleans, LA, pp. 1–8.
- Orange, A., James, P.W. and White, F.J. (2001). *Microchemical Methods for the Identification of Lichens*. British Lichen Society, London.
- Ørstedal, D.O. and Lewis Smith, R.I. (2001). *Lichens of Antarctica and South Georgia. A Guide to their Identification and Ecology*. Cambridge University Press, Cambridge.
- Stamatakis, A., Ludwig, T. and Meier, H. (2005). RAXML-III: a fast program for maximum likelihood – based inference of large phylogenetic trees. *Bioinformatics* **21**, 456–463.
- Stamatakis, A., Hoover, P. and Rougemont, J. (2008). A rapid bootstrap algorithm for the RAXML web-servers. *Systematic Biology* **57**, 758–771.
- Stenroos, S., Velmala, S., Pykälä, J. & Ahti, T. (2016). *Lichens of Finland*. Finnish Museum of Natural History Luomus and University of Helsinki, Helsinki.
- Swofford, D.L. (2002). *PAUP\* version 4.0 b10. Phylogenetic analysis using parsimony (\*and other methods)*. Sinauer, Sunderland, MA.
- Thomson, J.W. (1984). *American Arctic Lichens. 1. The Macrolichens*. Columbia University Press, New York.
- White, T.J., Bruns, T., Lee, S., and Taylor, J.W. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics, in M.A. Innis, D.H. Gelfand, J.J. Sninsky, & T.J. White (eds), *PCR Protocols: A guide to methods and applications*, pp. 315–322. Academic Press, New York.
- Zoller, S., Scheidegger, C. and Sperisen, C. (1999). PCR primers for the amplification of mitochondrial small subunit ribosomal DNA of lichen-forming ascomycetes. *Lichenologist* **31**, 511–516.